

New Tool for "Old" Nuclear Physics: FEL γ - Nucleus Colliders

H. Aktaş^a, N. Büget^b, A. K. Çiftçi^a, N. Meriç^a, S. Sultansoy^{b,c}, Ö. Yavaş^b

^a*Physics Department, Faculty of Sciences, Ankara University, 06100
Tandoğan, Ankara, TURKEY*

^b*Department of Engineering Physics, Faculty of Sciences, Ankara
University, 06100 Tandoğan, Ankara, TURKEY*

^c*Institute of Physics, Academy of Sciences, H.Cavid avenue 143, Baku,
AZERBAIJAN*

A new method has been considered to investigate the scattering reactions with photons on fully ionized nuclei. To induce γ -nucleus collisions a free electron laser and a heavy ion synchrotron have been considered bringing them together. Main parameters of the collider (especially LHC and HERA for the acceleration of nuclei) have been estimated. Rough calculations have also been made for the cross sections of the excited nuclei production. Finally, some design problems involving the collider have been considered.

I. INTRODUCTION

Linac-ring type ep and γp colliders have been proposed recently as the fourth way to reach energies in the TeV scale [1]. If future linear e^+e^- colliders (or special e -linacs) are constructed near the proton rings of HERA (ep collider, Germany) and FNAL ($\bar{p}p$ collider, USA) or LHC (pp collider, CERN) under construction, a number of additional opportunities will arise. For example [2],

$$\begin{aligned} \text{LHC} \otimes \text{TESLA} = & \text{LHC} \oplus \text{TESLA} \\ & \oplus \text{TeV scale } ep \text{ collider [3] [4] [5] [6]} \\ & \oplus \text{TeV scale } \gamma p \text{ collider [7] [8] [6]} \\ & \oplus \text{Multi-TeV scale } e - \text{nucleus and } \gamma - \text{nucleus colliders [9] [6]} \\ & \oplus \text{FEL } \gamma - \text{nucleus collider} \end{aligned}$$

In this paper we deal with the last collider. As known, TESLA can operate as Free Electron Laser (FEL) in the X-ray region [10]. Collision of the FEL beam with the bunch of nuclei may give a unique possibility to investigate "old" nuclear phenomena in rather unusual conditions. In fact, keV energy FEL photons will be seen as a "laser" beam in the MeV energy range in the rest frame of the nucleus. Earlier investigations on this subject involve bremsstrahlung photons having continuous energy spectrum. Moreover, since the accelerated nuclei are fully ionized, the background which may be induced by the low-shell electrons will be eliminated. This option needs further investigations from both the accelerator and the nuclear physics point of view.

In section II general considerations of the proposed collider have been presented. Luminosities of γ -nucleus collisions have been estimated in section III. Next, in order to illustrate the physics search capacity of the proposed machines two processes have been given. Finally, some concluding remarks have been presented in section V.

II. GENERAL CONSIDERATIONS

The main elements of the proposed machine are illustrated in Fig.1. The γ -beam produced by the Free Electron Laser based on TESLA (or a special e -linac) collides with the nucleus beam from HERA or LHC. In the nucleus rest frame the energy of FEL photon is multiplied by the Lorentz factor γ_N of the nucleus ($\gamma_N \simeq 500$ for HERA and $\gamma_N \simeq 3000$ for LHC). The FEL photon wavelength is given by [10]

$$\lambda = \frac{l_u}{2\gamma_e^2} \left(1 + \frac{k^2}{2} \right), \quad (1)$$

where l_u is the period length of an undulator, $\gamma_e = E_e/m_e$ is the Lorentz factor of the electron (E_e and m_e being the energy and mass of the electron), $k = eB_u l_u / 2\pi m_e$ is the undulator parameter in which $e = \sqrt{4\pi\alpha}$ (α being the fine structure constant) and B_u is the peak value of the magnetic field in the undulator. For illustration, let us use the design parameters [10] of the TESLA Test Facility (TTF) FEL, which are presented in Table 1. The nucleus will "see" TTF FEL beam as photons with an energy of 96.5 keV at HERA and 579 keV at LHC. Last values may be adjusted by changing E_e , B_u , l_u or γ_N . With the necessary adjustment the energy region can be made suitable for the investigation of the electromagnetic excitations of the nucleus.

The excited nucleus will turn to the ground state at a distance $l = \gamma_N \times \tau_N \times c$ from the collision point, where τ_N is the lifetime of the excited state in the nucleus rest frame and c is the speed of light. As an example, for the 4847.2 keV excitation of ^{208}Pb nucleus at LHC (which will be considered in section IV) $l = 6 \times 10^{-5}$ m. Therefore, the detector should be placed close to the collision region. The 5 MeV energy photons emitted in the rest frame of the nucleus will be seen in the detector as high energy photons with energies up to 30 GeV.

III. LUMINOSITY ESTIMATIONS

In this section we present rough estimations of FEL γ - nucleus luminosities for HERA and LHC based machines. Luminosity of the γ -nucleus collision is given by

$$L = \frac{n_\gamma n_{nuc}}{s_{eff}} f, \quad (2)$$

where n_γ and n_{nuc} are the number of photons and nuclei in the FEL and nucleus bunches, respectively, $f = n_b \times f_{rep}$ is the collision frequency (in which n_b is the number of electron bunches per pulse and f_{rep} is the repetition rate) and $s_{eff} = 4\pi\sigma_x^{eff}\sigma_y^{eff}$. In the last expression σ 's denote the size of the beam on the coordinate axis. σ_x^{eff} and σ_y^{eff} , on the other hand, are the values which are obtained by choosing the bigger ones of the corresponding horizontal and vertical sizes of the FEL and the nucleus beams.

Another parameter which is important in the detector design is the luminosity per collision which is given by

$$L_0 = \frac{n_\gamma n_{nuc}}{s_{eff}}. \quad (3)$$

Of course, the bunch structure of linac pulses should be chosen in accordance with the structure of the nucleus beam.

In the estimations given below we have used HERA ^{12}C and LHC ^{208}Pb beam parameters [6], which are presented in Table 2. Substituting the corresponding values from Tables 1 and 2 into Eqs. (2) and (3) luminosity values given in Table 3 have been obtained. The values $L = 5 \times 10^{31} cm^{-2} s^{-1}$ for HERA (^{12}C) and $L = 10^{30} cm^{-2} s^{-1}$ for LHC (^{208}Pb) based FEL γ - nucleus colliders seems to be quite realistic estimations.

IV. PROCESS EXAMPLES

The cross section for the resonant photon scattering is given by the well-known Breit-Wigner formula:

$$\sigma(\gamma, \gamma') = \frac{\pi}{E^2} \times \frac{2J_{exc} + 1}{2 \times (2J_0 + 1)} \times \frac{B_{in} B_{out} \Gamma^2}{(E - E_R)^2 + \Gamma^2/4} \quad (4)$$

where E is the c.m. energy of the incoming photon (in our case it is very close to that in the rest frame of the nucleus), J_{exc} and J_0 are spins of the excited and ground states of the nucleus, B_{in} and B_{out} are the branching fractions of the excited nucleus into the entrance and exit channels, respectively, E_R is the energy at the resonance and Γ is the total width of the excited nucleus. The number of the scattering events from the nuclei of ^{12}C and ^{208}Pb have roughly been calculated below taking their excited states of 4438 keV [11] and 4842.2 keV [12] [13], respectively. One can easily obtain corresponding values for other excitations and other nuclei using appropriate informations from [14].

A. E = 4438 KeV Excitation of ^{12}C at FEL γ - HERA

In this case FEL photons with $\omega_0 = 8876$ eV are needed. According to Eq. (1) this corresponds to $E_e = 6.78$ GeV if other parameters of the undulator are fixed. Using the values of $J_{exc} = 2$, $J_0 = 0$, $\Gamma \simeq 10.8 \times 10^{-3}$ eV, $B_{in} = B_{out} = 1$, the resonant scattering cross-section has been obtained by Eq. (4) as

$$\sigma^{res} \simeq 6.2 \times 10^{-22} cm^2. \quad (5)$$

Taking into account the energy spread of FEL and the nucleus beams ($\Delta E_\gamma/E_\gamma = \Delta\omega/\omega_0 = 10^{-3}$ and $\Delta E_C/E_C = 10^{-4}$), the approximate value of averaged cross-section has been found to be

$$\sigma^{av} \simeq \sigma^{res} \frac{\Gamma}{\Delta E_\gamma} \simeq 1.5 \times 10^{-27} cm^2 \quad (6)$$

where $\Delta E_\gamma = 4438$ eV. This value corresponds to 6.4×10^9 events per day for $L = 5 \times 10^{31} cm^{-2} s^{-1}$.

B. E = 4842.2 KeV Excitation of ^{208}Pb at FEL γ - LHC

In this case $\omega_0 = 1616$ eV and $E_e = 2.89$ GeV. Substituting $J_{exc} = 1$, $J_0 = 0$, $\Gamma \simeq 5$ eV and $B_{in} = B_{out} = 1$ into Eq. (4) the resonant cross-section has been estimated as

$$\sigma^{res} \simeq 3.1 \times 10^{-22} cm^2 \quad (7)$$

and, consequently, with the same energy spreads given above, the average cross-section as

$$\sigma^{av} \simeq 0.3 \times 10^{-24} cm^2$$

which corresponds to 2.5×10^{10} events per day for $L = 10^{30} cm^{-2} s^{-1}$.

V. CONCLUSION

In this study a new method has been presented to investigate the nuclear excitations. Although the quantitative description was given for only two nuclei it can easily be extended for others too. The method has the advantage that the accelerated ionized nuclei see the keV energy FEL photons as a laser beam in the MeV energy range. In principle, FEL γ beam can be obtained from existing HERA and LEP ring electron beams, too. This option will be considered elsewhere.

For future investigations the determination of the following nuclear quantities might be promising: scattering cross-sections, spins and parity assignments of the excited states.

ACKNOWLEDGMENTS

This work is partially supported by Turkish State Planning Organization under Grant No 97.K.120420.

-
- [1] Proceedings of the First International Workshop on Linac-Ring Type ep and γ p Colliders, 9-11 April 1997, Ankara, Turkey. To be published in Turkish Journal of Physics.
 - [2] S. Sultansoy "Four Ways to TeV Scale" in Ref.[1].
 - [3] M. Tigner, B. Wiik and F. Willeke, Proc. 1991 IEEE Particle Accelerator Conference, Vol.5, p.2910.
 - [4] Z. Z. Aydin, A. K. Çiftçi and S. Sultansoy, Nucl. Instrum. Meth. A351 (1994) 261.
 - [5] R. Brinkmann and M. Dohlus, DESY preprint M-95-11, Hamburg (1995).
 - [6] R. Brinkmann et al., DESY preprint 97-239, Hamburg (1997).
 - [7] S. F. Sultanov, ICTP preprint IC/89/409, Trieste (1989).
 - [8] A. K. Çiftçi, S. Sultansoy, Ş. Türköz and Ö. Yavaş, Nucl. Instrum. Meth. A365 (1995) 317.
 - [9] Z. Z. Aydin et al., Proceedings of the 28'th Int. Conf. on High Energy Physics, eds Z. Ajduk and A. K. Wroblewski, World Scientific, Singapore, 1996, p.1752.
 - [10] TESLA-FEL 95-03, June 1995, A VUV Free Electron Laser at The TESLA Test Facility at DESY.
 - [11] K. P. Schelhaas et al., Nucl. Phys. A506 (1990) 307.
 - [12] C. P. Swann, Phys. Rev. Lett. 32 (1974) 1449.
 - [13] R. M. Laszewski and P. Axel, Phys. Rev. C 19 (1979) 342.
 - [14] Nuclear Data Sheets, Academic Press, Inc.

Table 1. TTF FEL parameters

Electron beam energy E_e , GeV	1
Number of electrons per bunch n_e , 10^{10}	0.624
Pulse length, μs	1000
Number of bunches per pulse n_b	7200
Repetition rate, Hz	10
Period length of an undulator l_u , mm	27.3
Peak field in the undulator B_u , T	0.497
Undulator parameter k	1.3
Photon wavelength λ , nm	6.4
Photon energy ω_0 , eV	193
Number of photons per bunch n_γ , 10^{13}	4
Energy spread of photons $\Delta\omega/\omega_0$, 10^{-3}	1
rms beam size $\sigma_{x,y}$, μm	50

Table 2. Parameters of nucleus beams

	^{12}C at HERA	^{208}Pb at LHC
Maximum beam energy, TeV	4.9	574
Particles per bunch n_{nuc} , 10^8	80	0.94
Normalized emittance ε^N , mm $\times$ mrاد	1.25	1.4
Amplitude function at IP β^* , cm	20	20
rms beam size at IP $\sigma_{x,y}$, μm	22	9.7
Bunch spacing, ns	192	100
Number of bunches in FEL pulse n_b	5208	10000

Table 3. Luminosities of FEL γ - nucleus collisions

	HERA ^{12}C	LHC ^{208}Pb
Luminosity L , $10^{30} \text{ cm}^{-2} \text{ s}^{-1}$	53	1.1
Luminosity per collision L_0 , 10^{26} cm^{-2}	7.2	0.17




Fig.1: General schematic view of the proposed design